

Structure and texture of the quark mass matrix

Maritza de Coss^a and Rodrigo Huerta^b

^aDepartamento de Física Aplicada,
Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional,
Unidad Mérida, A. P. 73 Cordemex 97310, Mérida, Yucatán, México.

Starting from a weak basis in which the up (or down) quark matrix is diagonal, we obtain an exact set of equations for the quark mass matrix elements in terms of known observables. We make a numerical analysis of the down (up) quark mass matrix. Using the data available for the quark masses and mixing angles at different energy scales, we found a numerical expression for these matrices. We suggest that it is not possible to have an specific texture from this analysis. We also examine the most general case when the complex phases are introduced in the mass matrix. We find the numerical value for these phases as a function of δ , the CP-violation phase.

PACS number(s): 12.15.Ff

1. Introduction

The known observables in the Yukawa sector of the Standard Model (SM) are six quark mass eigenvalues plus the four parameters of the Cabibbo-Kobayashi-Maskawa (CKM) mixing matrix. These parameters are related to the quark mixing matrices through the diagonalization procedure. The two quark mass matrices with three generations depend on 36 parameters in general (18 each). There are many possibilities in the literature that propose different structures for these matrices. In particular, there are studies considering the mass matrix as hermitian, symmetric [2–4] or non-hermitian [5]. To reduce the number of parameters, zeros are introduced in the mass matrices leading to the terminology of texture [4]. Depending on the particular texture it is possible to find relations among masses and mixing angles. It would be of interest to explore what is the texture at which the present data is pointing.

In this work we obtain a set of exact equations relating the mass matrix parameters and the known observables. We perform a numerical analysis with those algebraic equations, using the available data, and we find the structure of the quark mass matrices. We do this analysis for different energy scales using the renormalization group equation to find the evolution of the known observables. In the next section we obtain the exact set of equations from which we observe the relation among the measured quantities and the quark mass matrix elements, including the phases of the latter. In section III, we find the evolution of these matrix elements. From this analysis we find their hierarchy. Finally, in section IV we summarize our results.

2. Expressions for the mass matrix elements

We start from the terms of the mass and the weak charged-current of the standard model Lagrangian which are important to us,

$$\mathcal{L}_{mass} = \bar{u}'_L \mathbf{m}_u u'_R + \bar{d}'_L \mathbf{m}_d d'_R + g_L \bar{u}'_L \mathbf{W}_L^\dagger d'_L + c.h., \quad (1)$$

where the quark mass matrices are general complex matrices whose size depends on the number of generations. We can rotate the weak states to get a diagonal mass matrix by making the usual transformations [1],

$$\begin{aligned} \mathcal{L}_{mass} &= \bar{u}'_L \mathbf{R}_L^u (\mathbf{R}_L^u)^\dagger \mathbf{m}_u (\mathbf{S}_R^u)^\dagger \mathbf{S}_R^u u'_R + \bar{d}'_L \mathbf{R}_L^d (\mathbf{R}_L^d)^\dagger \mathbf{m}_d (\mathbf{S}_R^d)^\dagger \mathbf{S}_R^d d'_R \\ &\quad + g_L \bar{u}_L (\mathbf{R}_L^u)^\dagger \mathbf{W}_L^\dagger \mathbf{R}_L^d d_L + c.h. \\ &= \bar{u}_L \mathbf{M}_u u_R + \bar{d}_L \mathbf{M}_d d_R + g_L \bar{u}_L \mathbf{W}_L^\dagger \mathbf{V}_{CKM} d_L + c.h. \end{aligned} \quad (2)$$

where the unitary matrices $\mathbf{R}_L^{u,d}$ and $\mathbf{S}_R^{u,d}$ rotate the weak basis to the mass eigenstates,

$$\begin{aligned} u'_L &= \mathbf{R}_L^u u_L, & d'_L &= \mathbf{R}_L^d d_L \\ u'_R &= \mathbf{S}_R^u u_R, & d'_R &= \mathbf{S}_R^d d_R. \end{aligned} \quad (3)$$

In this way we obtain the biunitary transformations which transform the mass matrices $\mathbf{m}_u$ and $\mathbf{m}_d$ to their diagonal form,

$$\begin{aligned} \mathbf{M}_u &= (\mathbf{R}_L^u)^\dagger \mathbf{m}_u \mathbf{S}_R^u, \\ \mathbf{M}_d &= (\mathbf{R}_L^d)^\dagger \mathbf{m}_d \mathbf{S}_R^d. \end{aligned} \quad (4)$$

The CKM mixing matrix [6] is then given by

$$\mathbf{V}_{CKM}^L = (\mathbf{R}_L^u)^\dagger \mathbf{R}_L^d. \quad (5)$$

In the case of having a charged right-handed current of the type $g_R \bar{u}'_R \mathbf{W}_R^\dagger d'_R$ in the Lagrangian [7] (neglecting the possible mixing between the gauge bosons), we would have

$$\mathbf{V}_{CKM}^R = (\mathbf{S}_R^u)^\dagger \mathbf{S}_R^d. \quad (6)$$

Without any loss of generality we can have one of the mass matrices diagonal [8,9]. Assume first that

$$\mathbf{m}_u = \begin{pmatrix} m_u & 0 & 0 \\ 0 & m_c & 0 \\ 0 & 0 & m_t \end{pmatrix}, \quad (7)$$

then the matrices $\mathbf{R}_L^u$ and $\mathbf{S}_R^u$ are equal to the unit matrix. From equations (5) and (6) we obtain

$$\begin{aligned} \mathbf{V}_{CKM}^L &= \mathbf{R}_L^d, \\ \mathbf{V}_{CKM}^R &= \mathbf{S}_R^d. \end{aligned} \quad (8)$$

In this case the diagonal mass matrix d is given by,

$$\mathbf{M}_d = (\mathbf{V}_{CKM}^L)^\dagger \mathbf{m}_d \mathbf{V}_{CKM}^R. \quad (9)$$

One can make $\mathbf{V}_{CKM}^R$ equal to $\mathbf{V}_{CKM}^L$ [8] and g_L equal to g_R assuming *manifest* left-right symmetry in the flavor sector, and then one can see that we need only to consider a single mixing matrix which is responsible of the diagonalization,

$$\mathbf{M}_d = (\mathbf{V}_{CKM})^\dagger \mathbf{m}_d \mathbf{V}_{CKM}. \quad (10)$$

This is equivalent to assuming that $\mathbf{m}_d$ is hermitian. Now we can express the mixing angles in terms of the mass eigenvalues m_i and the mass matrix elements m_{ij} . To this end we take an hermitian mass matrix and choose the standard form for $\mathbf{V}_{CKM}$,

$$\mathbf{V}_{CKM} = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta} & c_{23}c_{13} \end{pmatrix}, \quad (11)$$

so we have

$$\mathbf{V}_{CKM} \begin{pmatrix} m_1 & 0 & 0 \\ 0 & m_2 & 0 \\ 0 & 0 & m_3 \end{pmatrix} = \begin{pmatrix} m_{11} & m_{12}e^{i\delta_{12}} & m_{13}e^{i\delta_{13}} \\ m_{12}e^{-i\delta_{12}} & m_{22} & m_{23}e^{i\delta_{23}} \\ m_{13}e^{-i\delta_{13}} & m_{23}e^{-i\delta_{23}} & m_{33} \end{pmatrix} \mathbf{V}_{CKM}. \quad (12)$$

Considering the magnitudes of the elements (2,3), (1,3) and (1,2) on both sides of this equation we get the mixing angles in exact form [10],

$$\frac{s_{23}}{c_{23}} = \frac{(m_3 - m_{11})m_{23} + m_{13}m_{12}}{(m_3 - m_{22})(m_3 - m_{11}) - m_{12}^2} \quad (13)$$

$$\frac{s_{13}}{c_{13}} = \frac{m_{12}s_{23} + m_{13}c_{23}}{m_3 - m_{11}} \quad (14)$$

$$\frac{s_{12}}{c_{12}} = \frac{m_{12}c_{23} - m_{13}s_{23}}{(m_2 - m_{11})c_{13} + (m_{12}s_{23} + m_{13}c_{23})s_{13}} \quad (15)$$

where $s_{ij} = \sin \theta_{ij}$, $c_{ij} = \cos \theta_{ij}$. To obtain the quark masses as functions of the matrix elements we use the fact that $\mathbf{m}_d$ satisfies the following characteristic equation,

$$\begin{aligned} \det(\mathbf{m}_d - m\mathbf{1}) &= -m^3 + (m_{11} + m_{22} + m_{33})m^2 - (m_{11}m_{22} + m_{11}m_{33} \\ &\quad + m_{22}m_{33} - m_{23}^2 - m_{13}^2 - m_{12}^2)m + m_{11}(m_{22}m_{33} - m_{23}^2) \\ &\quad - m_{12}(m_{12}m_{33} - m_{13}m_{23}) + m_{13}(m_{12}m_{23} - m_{13}m_{22}) \\ &= 0. \end{aligned} \quad (16)$$

The eigenvalues m_i also satisfy the equation

$$\begin{aligned} (m_1 - m)(m_2 - m)(m_3 - m) &= -m^3 + (m_1 + m_2 + m_3)m^2 \\ &\quad - (m_1m_2 + m_1m_3 + m_2m_3)m + m_1m_2m_3 \\ &= 0. \end{aligned} \quad (17)$$

After equating the coefficients with the same power of m in (16) and (17) we get,

$$m_1 + m_2 + m_3 = m_{11} + m_{22} + m_{33} \quad (18)$$

$$m_1 m_2 + m_1 m_3 + m_2 m_3 = m_{11} m_{22} + m_{11} m_{33} + m_{22} m_{33} - m_{23}^2 - m_{13}^2 - m_{12}^2 \quad (19)$$

$$m_1 m_2 m_3 = m_{11}(m_{22} m_{33} - m_{23}^2) - m_{12}(m_{12} m_{33} - m_{13} m_{23}) + m_{13}(m_{12} m_{23} - m_{13} m_{22}). \quad (20)$$

Going back to eq. (12), we compare the phases of elements (1,2), (1,3) and (2,3) on both sides of the equation to get the relations,

$$\tan(\delta_{12}) = \frac{-(m_3 - s_{12}^2 m_2 - m_1) s_{13} s_{23} \sin \delta}{(m_2 - m_1) s_{12} + (m_3 - s_{12}^2 m_2 - m_1) s_{13} s_{23} \cos \delta} \quad (21)$$

$$\tan(\delta_{13}) = \frac{(m_3 - s_{12}^2 m_2 - m_1) s_{13} \sin \delta}{(m_2 - m_1) s_{12} s_{23} - (m_3 - s_{12}^2 m_2 - m_1) s_{13} \cos \delta} \quad (22)$$

$$\tan(\delta_{23}) = \frac{(m_2 - m_1)(1 + s_{23}^2) s_{12} s_{13} \sin \delta}{[m_3 - (s_{12}^2 - s_{13}^2) m_1 - (1 - s_{12}^2 s_{13}^2) m_2] s_{23} - (m_2 - m_1)(1 - s_{23}^2) s_{12} s_{13} \cos \delta}. \quad (23)$$

We observe that the phases δ_{ij} are given in terms of known observables and that they are independent of the m_{ij} . We consider important to remark that the set of equations (13-15), (18-20) and (21-23) are exact relations among quark mass matrix elements and known observables.

3. Structure and evolution of the mass matrix elements

The structure and evolution of the mass matrix can be found by numerically solving the set of equations obtained in the previous section. To this end we give in the following the quark mass eigenvalues, the mixing angles and the δ -phase at different scales. For the quark mass eigenvalues we use the quantities shown in Table 1 [11].

We use the following values for the mixing angles at low energy scales (1 GeV to m_t) [12]

$$\begin{aligned} \sin \theta_{12} &= 0.2225 \pm 0.0021 \\ \sin \theta_{23} &= 0.04 \pm 0.0018 \\ \sin \theta_{13} &= 0.0035 \pm 0.0009 \end{aligned} \quad (24)$$

and for the CP-phase we take [13],

$$\delta = (66.5 \pm 30.5)^\circ. \quad (25)$$

To calculate the mixing angles at $10^9 GeV$ and M_X , where M_X is the unification scale of SUSY ($M_X = 2 \times 10^{16} GeV$), we use the formalism in ref. [14]. In this work Kielanowski *et al.*, find the energy dependence of the $|V_{ij}|$. Recalling that from eq. (11), the mixing angles are related to the magnitudes of the V_{ij} matrix elements. In particular we are

Table 1
Running quark masses (in units of GeV).

<i>Scale</i>	<i>u</i>	<i>d</i>	<i>s</i>
<i>1 GeV</i>	0.00488 ± 0.00057	0.00981 ± 0.00065	0.1954 ± 0.0125
m_Z	0.00233 ± 0.00045	0.00469 ± 0.00066	0.0934 ± 0.0130
m_t	0.00223 ± 0.00043	0.00449 ± 0.00064	0.0894 ± 0.0125
$10^9 GeV$	0.00128 ± 0.00025	0.00260 ± 0.00037	0.0519 ± 0.072
M_X	0.00094 ± 0.00018	0.00194 ± 0.00028	0.0387 ± 0.054
	<i>c</i>	<i>b</i>	<i>t</i>
<i>1 GeV</i>	1.506 ± 0.048	7.18 ± 0.59	475 ± 86
m_Z	0.677 ± 0.061	3.00 ± 0.11	181 ± 13
m_t	0.646 ± 0.059	2.85 ± 0.11	171 ± 12
$10^9 GeV$	0.371 ± 0.033	1.51 ± 0.06	109 ± 16
M_X	0.272 ± 0.024	1.07 ± 0.04	84 ± 18

interested in the magnitudes of $|V_{us}|$, $|V_{cb}|$ and $|V_{ub}|$. The evolution of these magnitudes are found to be given by,

$$|V_{ub}|^2 = \frac{|V_{ub}^0|^2}{|V_{tb}^0|^2[h(t)^2 - 1] + 1} \quad (26)$$

$$|V_{cb}|^2 = \frac{|V_{cb}^0|^2}{|V_{tb}^0|^2[h(t)^2 - 1] + 1} \quad (27)$$

where $|V_{ij}^0|$ are the initial values of the CKM matrix elements and $h(t)$ is expressed as

$$h(t) = \left(\frac{1}{1 - \frac{3(b+2)}{(4\pi)^2} m_t^2(t_0) \int_{t_0}^t r_g(\tau) d\tau} \right)^{\frac{c}{2(b+2)}}$$

with

$$r_g(t) = \exp \left(\frac{2}{(4\pi)^2} \int_{t_0}^t \alpha_1^u(\tau) d\tau \right).$$

In the SM we have,

$$\alpha_1^u(t) = - \left(\frac{17}{20} g_1^2 + \frac{9}{4} g_2^2 + 8 g_3^2 \right).$$

with

$$g_i(t) = \frac{g_i(t_0)}{\sqrt{1 - \frac{2b_i g_i^2(t_0)(t-t_0)}{(4\pi)^2}}},$$

and $t = \ln(E/m_t)$. $|V_{us}|$ is obtained from unitarity. We evaluate at $t_1 = \ln(\frac{10^9}{171}) \sim 15.5816$ and $t_2 = \ln(\frac{2 \times 10^{16}}{171}) \sim 32.3928$. We assume there is no running for the δ phase [15]. We can collect all the numerical results in Table 2.

Table 2
Elements of mass matrix d

Scale	m_{11}	$m_{12}e^{i\delta_{12}}$	$m_{13}e^{i\delta_{13}}$
1 GeV	$.0194 \pm .0030$	$(.0412 \pm .0023)e^{-i(1.26 \pm .5)^\circ}$	$(.0235 \pm .0066)e^{-i(70.05 \pm 31.22)^\circ}$
m_Z	$.0094 \pm .0020$	$(.0196 \pm .0019)e^{-i(1.11 \pm .43)^\circ}$	$(.0097 \pm .0027)e^{-i(70.58 \pm 31.31)^\circ}$
m_t	$.0090 \pm .0019$	$(.0188 \pm .0018)e^{-i(1.10 \pm .42)^\circ}$	$(.0092 \pm .0026)e^{-i(70.61 \pm 31.32)^\circ}$
$10^9 GeV$	$.0052 \pm .0010$	$(.0110 \pm .0009)e^{-i(1.31 \pm .49)^\circ}$	$(.0055 \pm .0015)e^{-i(71.04 \pm 31.38)^\circ}$
M_X	$.0039 \pm .0008$	$(.0082 \pm .0007)e^{-i(15.89 \pm 6.7)^\circ}$	$(.0048 \pm .0013)e^{-i(66.95 \pm 30.6)^\circ}$
	m_{22}	$m_{23}e^{i\delta_{23}}$	m_{33}
1 GeV	$.1987 \pm .0309$	$(.2798 \pm .0193)e^{i(.03 \pm .01)^\circ}$	$(7.167 \pm .6089)$
m_Z	$.0936 \pm .0161$	$(.1163 \pm .0061)e^{i(.03 \pm .01)^\circ}$	$(2.995 \pm .1155)$
m_t	$.0896 \pm .0155$	$(.1105 \pm .0060)e^{i(.03 \pm .01)^\circ}$	$(2.845 \pm .1155)$
$10^9 GeV$	$.0525 \pm .0081$	$(.0671 \pm .0035)e^{i(.03 \pm .01)^\circ}$	$(1.507 \pm .0529)$
$M_X GeV$	$.0400 \pm .0062$	$(.0571 \pm .0030)e^{i(.04 \pm .01)^\circ}$	$(1.067 \pm .0424)$

Table 3
Elements of mass matrix u

Scale	m_{11}	$m_{12}e^{i\delta_{12}}$	$m_{13}e^{i\delta_{13}}$
1 GeV	$.0887 \pm .0248$	$(.3923 \pm .0216)e^{-i(9.60 \pm 4.12)^\circ}$	$(1.662 \pm .4580)e^{-i(66.92 \pm 30.60)^\circ}$
m_Z	$.0387 \pm .0074$	$(.1716 \pm .0119)e^{-i(8.25 \pm 3.32)^\circ}$	$(.6284 \pm .1657)e^{-i(67.99 \pm 30.61)^\circ}$
m_t	$.0369 \pm .0070$	$(.1635 \pm .0114)e^{-i(8.18 \pm 3.29)^\circ}$	$(.5936 \pm .1564)e^{-i(67.0 \pm 30.61)^\circ}$
$10^9 GeV$	$.0220 \pm .0053$	$(.1001 \pm .0075)e^{-i(11.53 \pm 4.85)^\circ}$	$(.4340 \pm .1138)e^{-i(67.0 \pm 30.60)^\circ}$
M_X	$.0171 \pm .0050$	$(.0816 \pm .0071)e^{-i(54.1 \pm 26.6)^\circ}$	$(.4107 \pm .1078)e^{-i(66.54 \pm 30.51)^\circ}$
	m_{22}	$m_{23}e^{i\delta_{23}}$	m_{33}
1 GeV	$2.266 \pm .563$	$(18.96 \pm 2.04)e^{i(.0032 \pm .0013)^\circ}$	474 ± 86
m_Z	$.937 \pm .118$	$(7.22 \pm .47)e^{i(.0038 \pm .0014)^\circ}$	181 ± 13
m_t	$.891 \pm .118$	$(6.82 \pm .44)e^{i(.0039 \pm .0014)^\circ}$	171 ± 12
$10^9 GeV$	$.591 \pm .104$	$(5.0 \pm .43)e^{i(.0035 \pm .0013)^\circ}$	109 ± 13
M_X	$.529 \pm .116$	$(4.64 \pm .48)e^{i(.0336 \pm .0138)^\circ}$	84 ± 13

If we consider m_d to be diagonal, instead of m_u , a similar analysis determines the structure of the quark mass matrix m_u . The result of this analysis is shown in Table 3.

From these numerical expressions for the mass matrices, one can see that *no* element is consistent with zero at 3σ , for any given scale. We notice the following hierarchy among the magnitudes of the mass matrix elements,

$$m_{11} \approx m_{13} < m_{12} < m_{22} < m_{23} \ll m_{33}. \quad (28)$$

This hierarchy is similar to the one obtained by Fritzsch [16]: $m_{11}, m_{12}, m_{13} \ll m_{23}, m_{22} \ll m_{33}$. We also have a hierarchy among the phases

$$\delta_{23} \ll \delta_{12} \ll \delta_{13} \quad (29)$$

at low energy scales. At SUSY scales this hierarchy is no longer valid and we have instead $\delta_{23} \ll \delta_{12} \approx \delta_{13}$.

4. Conclusions

To summarize, starting with the weak basis, for which one of the quark mass matrices is diagonal, we find exact relations that are analysed numerically. We obtain numerical expressions at different scales, using the mixing angles and quark masses as input data. From this analysis we conclude that *no* texture is from 1 GeV up to $2 \times 10^{16}\text{ GeV}$ at 3σ . We also find the explicit dependence of δ_{ij} in terms of the quark masses and the CKM mixing angles. Numerical evaluation shows that we have $\delta_{23} \ll \delta_{12} \ll \delta_{13}$ at low scales and $\delta_{23} \ll \delta_{12} \sim \delta_{13}$ at SUSY scales. We expect in the near future better measurements for the known observables in the Yukawa sector of the SM. When the measurements of these observables becomes more precise, we will be able to draw more definitive conclusions about the existence of a particular texture in the quark mass matrices.

We would like to thank useful conversations with A. Bouzas, B. Desai, E. Ma, G. Sanchez-Colón and J. Wudka. One of us (RH) wants to thank the hospitality of the Department of Physics at UCR, where part of this work was done. This work was partially supported by Conacyt (México).

REFERENCES

1. S. Weinberg, The Quantum Theory of Fields (Vol. II), Cambridge University Press, 1996.
2. H. Fritzsch, Phys. Lett. B 70 (1977) 436; 73 (1978) 317.
3. H. Georgi and C. Jarlskog, Phys. Lett. B 86 (1979) 297.
4. P. Ramond, R. G. Roberts and G.G. Ross, Nucl. Phys. B406 (1993) 19.
5. G. C. Branco and J. I. Silva-Marcos, Phys. Lett. B 331 (1994) 390.
6. N. Cabibbo, Phys. Rev. Lett. 10 (1963) 531; M. Kobayashi and T. Maskawa, Prog. Theor. Phys. 49 (1973) 652.
7. R. N. Mohapatra, Unification and Supersymmetry: The Frontiers of Quark-Lepton (2nd. edition), Springer Verlag, 1992.
8. E. Ma, Phys. Rev. D 43 (1991) 2761.
9. Y. Koide, H. Fusaoka and C. Habe, Phys. Rev. D 46 (1992) 4813.
10. A. Rašin, Phys. Rev. D 58 (1998) 96012.
11. H. Fusaoka and Y. Koide, Phys. Rev. D 57 3986 (1998) 3986.
12. D. E. Groom et al., Particle Data Group, Eur. Phys. J. C15 (2000) 1, p. 103.
13. A. Ali and D. London, Eur. Phys. J. C 9 (1999) 687. They quote $36^\circ \leq \delta \leq 97^\circ$.
14. P. Kielanowski, S.R. Juárez, J. C. Mora, Phys. Lett. B (2000) 179.
15. C. Balzereit *et al.*, Eur. Phys. J. C 9 (1999) 197.
16. H. Fritzsch and Z. Xing, Phys. Lett. B 413 (1997) 396.